



Booster

w/v 120g/L N 120g/L Ca 30g/L Mg 10g/L B



NutriEssential products have the potential to improve yields, influence fruit and grain size, enhance colour and improve quality. NutriEssential activates plant mechanisms that impact the plant's natural defence mechanisms. NutriEssential products are formulated with balanced nutrition.

Crop	Rate/ha	Mini Dilution	Comments
Vegetables / Horticulture	3L-10L	1:100	Foliar: 3 applications prior to initial flower. Fertigation: Apply every 14-21 days from emergence or transplant.
Broadacre Crops - Winter or Summer; Irrigated or Rain-Fed	3L-10L	1:25	Apply as required from early tillering.
Vines	3L-10L	1:200	Foliar: Apply at clusters visible, flower buds separated and fruit set. Fertigation: Apply every 14-21 days as required.
Fruit, Nut & Other Tree Crops	3L-10L	1:300	Foliar: 3 applications from initial flower. Fertigation: Apply every 14-21 days.
Turf	100ml-200ml /100m2	1:10	Apply as required.

COMPATIBILITY: NutriEssential Booster is compatible with a wide range of commonly used agricultural chemicals, when mixing with other chemicals, always mix a small quantity by preforming a jar test and check for precipitate (fall out) in the mixture. If precipitation occurs, DO NOT proceed with application. If mixture does not precipitate and it's the first time you have used this product or combination of products on the target crop or the area, it's strongly recommended a small test area should be sprayed and observed.

NOTE: The suggested rates of application and dilutions should be used as a guide only. The rates of application and dilution necessary for optimum results may vary depending upon the user's particular environment and application processes. Advice on specific application rates should be sought from qualified agronomists. Where possible, it is recommended that regular tissue (leaf; fruit; grain) tests are conducted to determine actual plant nutrient status and availability during each growing cycle. Regular soil tests are recommended, preferably at least once per year. Avoided application under extreme weather conditions such as temperatures over 28C, high humidity, rapid drying conditions, frost, rain etc.

WARNING: Avoid ingesting product. Avoid contact with the eyes and skin. Wash hands after use. As farm management practices, including irrigation, application equipment and methods and grazing practices as well as environmental, climatic and soil conditions are beyond the control of Sonic Essentials Pty Ltd, Sonic Essentials Pty Ltd accepts no responsibility whatsoever for any direct or indirect loss or damage or other result following the use of this product whether used in accordance with directions or not, subject to any overriding statutory provision and provided that such liability under those provisions shall be limited to the replacement of the goods as supplied. This product is sold and the buyer accepts and uses this product subject to these conditions.

STORAGE & CARE OF EQUIPMENT: Store in a cool place, out of direct sunlight. Replace lid after use. This product can be corrosive to metals, in particular brass, copper, manganese and galvanized iron. Clean equipment after use and follow manufacturers maintenance advice.

Sonic Essentials Pty Ltd







